### **Drug Facts**

Uses' temporarily relieves minor aches and pains associated with arthritis

#### Warnings

Stop use and ask a doctor if symptoms persist for more than 7 days or worsen, new symptoms occur, or if redness or swelling is present. These could be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- Adults 18 years of age and older: At the onset of symptoms, dissolve 2 tablets under the tongue and repeat, if needed, every 15 minutes for 3 more doses. Then, dissolve 2 tablets under the tongue every hour until symptoms improve.
- Children under 18 years of age: Ask a doctor.

#### Other information

- do not use if glued carton end flaps are open or if the blister seal is broken
- store below 86°F (30°C)

# CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

"C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

# Drug Facts (continued) Inactive ingredients

croscarmellose sodium, lactose, magnesium stearate

## Questions or comments?

Arnicare.com BoironUSA.com Info@Boiron.com 1-800-B0IRON-1 (1-800-264-7661)

